

Graptolite Taxonomy and Classification

MU EN-ZHI



Mu En-zhi: Graptolite Taxonomy and Classification. *Bull. geol. Soc. Denmark*, vol. 35, pp. 203-207, Copenhagen, July 1st, 1987.

Graptolithina comprises chiefly six orders. Among them Graptoloidea and a part of Dendroidea known as Graptodendroids are planktonic in mode of life. Graptoloidea consists of three suborders namely Axonolipa, Axonocrypta and Axonophora. The families Dendrograptidae-Anisograptidae-Tetragraptidae and Didymograptidae-Isograptidae-Cardiograptidae-Diplograptidae-Monograptidae represent anagenetic grades. Some important evolutionary trends took place once again, representing cladogenetic divergences. All other families or subfamilies are offshoots of various grades. The suborder Axonocrypta is discussed in detail.

Mu En-zhi, Nanjing Institute of Geology and Palaeontology, Academia Sinica, Chimingssu, Nanjing, China, August 20th, 1986.

General Consideration

Graptolithina, a class of Hemichordata, comprises chiefly six orders known as Dendroidea, Graptoloidea, Tuboidea, Camaroidea, Stolonoidea and Crustoidea (Kozłowski, 1949, 1966; Bulman, 1970). The thecae in Dendroidea and Graptoloidea are regularly arranged in stipes, although Dendroidea has three kinds of thecae (autotheca, bitheca and stolotheca), while Graptoloidea has only one. In the graptoloid thecae, the proximal portion (protheca) and the distal portion (metatheca) are homologous with the stolotheca and autotheca respectively.

Among these orders, Dendroidea is the earliest in appearance (M. Cam.?, Late Cam.) and the latest in disappearance (Late Carb.); Graptoloidea ranged from Early Ordovician to Early Devonian; the other orders are rare in materials and known to occur from Ordovician to Silurian.

All the Tuboidea, Camaroidea, Stolonoidea, Crustoidea and most of Dendroidea are benthonic in the mode of life, whereas Graptoloidea and a part of Dendroidea are planktonic. The floating dendroids with free sicula appeared in the latest Cambrian (Fengshanian) and flourished in Early Xinchangian (Tremadocian), indicating a new stage in graptolite history. The graptodendroid family Anisograptidae comprising three subfamilies (quadriradiate Staurograp-

tinae, tri-radiate Anisograptinae and biradiate Adelograptinae) is derived from the floating *Diclyonema* due to the loss of dissepiments (Mu, 1974), while the reclined Psigraptidae Lin (1981) with isolated autothecae is an offshoot. Recently Zhao & Zhang (1985) proposed a new family Muenzhigraptidae with biform autothecae representing the direct ancestor of Psigraptidae.

Graptoloidea first appeared in the late Xinchangian (X_3) due to the loss of bithecae from Adelograptinae and flourished in early Ningkuoan (N_1), marking another new stage in graptolite history. Since then, Graptoloidea became the master of the quiet sea area in Ordovician and Silurian and even in Early Devonian.

Subdivision of the Order Graptoloidea

The subdivision of Graptoloidea has been treated by many graptolite workers. Two suborders, Axonolipa and Axonophora, proposed by Frech (1897) and emended by Ruedemann (1904, 1908) have been used for a long time (Mu, 1950; Obut, 1957; Mu, 1974; Yu & Fang, 1979). In fact, Axonolipa consists of Didymograptina and Dicellograptina of Lapworth (1880) and Axonophora consists of Diplograptina and Monograptina of Lapworth (1880). In the sixties, Jaanusson (1960) divided Graptoloidea into four suborders namely

Note: While the present volume was in press the editor received information on the death of Prof. Mu En-zhi.

Didymograptina, *Glossograptina*, *Diplograptina* and *Corynoidina*. Bulmann (1963) grouped *Corynoidina* into the suborder *Didymograptina* and divided the *Diplograptina* into two suborders, *Diplograptina* and *Monograptina*. Mu & Zhan (1966) established the suborder *Axonocrypta* in addition to *Axonolipa* and *Axonophora* based on the structure of the rhabdosome. Mu (1974) tentatively used *Didymograptina* and *Dicellograptina* as subdivisions of *Axonolipa* and *Diplograptina* and *Monograptina* as subdivisions of *Axonophora* respectively.

Axonolipa comprises the forms with nema free and the stipes pendent to reclined just like floating dendroids. Those in *Didymograptina* are developed in a primitive type. The initial bud originates from the porus on the ventral or dorsal side of the prosicula or the proximal part of metasicula. The proximal thecae grow downwards to nearly horizontally with one or two crossing canals, that is to say, the first or the second theca is dicalycal. The virgella is usually absent, or present in advanced forms, it begins at the proximal or middle part of the metasicula. The most primitive forms such as *Didymograptus* of *Didymograptidae*, *Tetragraptus* of *Tetragraptidae* and bitheca-lacking *Clonograptus* of *Dichograptidae* are the direct descendants of *Adelograptinae* of *Anisograptidae*. In *Dicellograptina*, the initial bud arises from the porus on the middle or distal part of metasicula. The proximal thecae grow horizontally to upwards with two or three crossing canals, that is to say, the second or third theca is dicalycal. The virgella is well developed, beginning usually at the proximal part of the metasicula. *Dicellograptus* of *Dicranograptidae* is the earliest derived from *Didymograptidae*.

In *Axonocrypta* the nema (virgula) is enclosed between the four to two scandent stipes. The sicula is relatively long. The initial bud originates from the porus on the prosicula or the proximal part of metasicula. It arises from the ventral, lateral or dorsal side of the sicula. The proximal thecae grow downwards or curved distally, with one or two crossing canals. The virgella is usually present in advanced form. Those with four or three stipes back to back (*Phyllograptidae*) are derived from *Tetragraptidae*, and those with two scandent stipes back to back (*Cardiograptidae*) or side to side (*Glossograptidae*) are derived

from *Isograptidae*, a descendant of *Didymograptidae* (and *Tetragraptidae*?).

The suborder *Axonophora* comprises those with virgula embedded in the median septum in biserial form (*Diplograptina*) or in the dorsal wall of the periderm in uniserial form (*Monograptina*). In *Diplograptina*, the initial bud originates from the porus on the distal part of metasicula and even near the sicular aperture. The proximal thecae in early forms are curved upwards with three or more crossing canals; in some later ones, the crossing canals increase throughout the rhabdosome without dicalycal theca (aseptate). The virgella is well developed, beginning at the proximal part of the metasicula even near the prosicula. The most primitive genus *Glyptograptus* of *Diplograptidae* is most probably derived from *Cardiograptidae* of *Axonocrypta* due to the fusion of the dorsal walls of two stipes, forming a median septum. *Exigraptus* seems to be a transitional form between *Cardiograptidae* and *Diplograptidae*. In China, the earliest species of *Glyptograptus*, *G. sinodentatus*, direct ancestor of *G. austrodentatus*, is more closely related to *Exigraptus* (Mu et al., 1979; Chen, 1982). It is questionable that *G. austrodentatus* is derived from *Maeandroglyptus* (Jenkins, 1980) and *G. dentatus* is related to *Phyllograptus elongatus* (Cooper & Fortey, 1983).

In *Monograptina* the initial bud originates from the sinus on the distal part of the metasicula near the sicular aperture. The proximal thecae grow directly upwards without crossing canal, i.e., no dicalycal theca. The earliest representatives of *Monograptina* are *Atavograptus* (Rickards, 1974), *Monoclimacis*? (Bjerreskov, 1975) and *Pristiograptus* (Li, in press) of *Monograptidae*, appearing abruptly in the beginning of Silurian (*persculptus* Zone) and representing another new stage of graptolite history. It is believed that *Monograptidae* is derived from *Diplograptidae* with discontinuity. *Peiragraptidae* and *Dimorphograptidae* are offshoots.

On the Suborder *Axonocrypta*

Axonocrypta linking with *Axonolipa* and *Axonophora* bears the fundamental characters between them in the structure of rhabdosome and

in the mode of development. There are three families in Axonocrypta, known as Phyllograptidae, Cardiograptidae and Glossograptidae (= Cryptograptidae).

Phyllograptidae is directly derived from Tetragraptidae of Axonolipa with *Tetragraptus phyllograptoides* and *Phyllograptus cor* as the transitional forms (Cooper & Fortey, 1982; Cooper & Lindholm 1985). The Family Phyllograptidae was revised by Hsu (1934) and Mu & Zhan (1966) to be composed of *Phyllograptus* and *Trigonograptus* (= *Pseudotrionograptus*) based on the structure of the rhabdosome. Recently, Cooper & Fortey (1982) revised this family to comprise *Phyllograptus* (s.s.) and a new genus *Xiphograptus* with two horizontal stipes, based on the origin of first theca from the dorsal side of the sicula and the presence of a virgella. In the writer's opinion, the orientation of the initial bud on the sicula is less taxonomic significance than the presence of virgella. Dorsal origin of initial bud on the sicula is also known in *Didymograptus rozkowskiae* and *Parazyograptus erraticus* (Kozłowski, 1954), but with no virgella. Since the presence of virgella is an advanced feature, the virgella-bearing *Phyllograptus* (s.s.) is more advanced than *Pseudophyllograptus* without a virgella, and the virgella-bearing *Xiphograptus* is an advanced member in Didymograptidae. It is noteworthy that the virgella-bearing *Pseudazyograptus incurvus* (Ekström), the type species of *Pseudazyograptus* (Mu et al., 1960), with elaborated thecae is an advanced member in Azyograptidae (Finney, 1980, p. 1199, Textfigs. 9, 10). True *Azyograptus* with simple thecae and without virgella occurs in Arenigian of Europe and Ningkuoan (N₄) of the Yangtze Region and the Jiangnan subregion of the S. China region in China, but is unknown in North America and Australasia.

In Phyllograptidae another advanced form is *Pseudotrionograptus uniformis* Mu et Lee, the type of species of *Pseudotrionograptus* (Mu and Lee, 1958, p. 417, PL. III, figs. 7–10). The detailed structure of the rhabdosome was described by Mu & Lee (1958) and illustrated by Hsu & Chen (1964, figs. 1, 4) based on the analysis of the pyritized thecae. The peculiar thecal characters and common canal and the parallel-sided rhabdosome in *Pseudotrionograptus uniformis*

are different from *Pseudotrionograptus ensiformis* as described by Rickards (1973, p. 600, figs. 1–3). They are not conspecific. Cooper has rightly distinguished these two species based on the materials of New Zealand (Cooper, 1979, p. 91, figs. 83b-c).

The pore of the common canal in *Pseudotrionograptus uniformis* corresponding to the "forenic foramina" in *Phyllograptus typus* illustrated by Cooper & Fortey (1982, figs. 71k, 74) is an advanced feature. The Spitzberg materials described by Fortey (1971) and Cooper & Fortey (1982) as *Pseudotrionograptus ensiformis* (4-stiped) and *P. minor* Mu et Lee (3-stiped) are possible new forms representing a genus intermediate between *Pseudophyllograptus* and *Pseudotrionograptus*, because the stipes are not entirely overlapped laterally as in *Pseudotrionograptus* and the serrated ventral margins of the stipes are clearly exposed. They are quite different from *Pseudotrionograptus ensiformis* and *P. minor* Mu et Lee. This genus seems to be identical with Cooper's Gen. 1 from New Zealand (Cooper, 1979, p. 93, figs. 84a-e). This genus is more primitive than *Pseudotrionograptus*, it occurs in a lower horizon. The other two families of Axonocrypta, Cardiograptidae and Glossograptidae with two scandent stipes back to back and side to side respectively are derived from isograptidae. Cardiograptidae with two stipes back to back (dipleural) developed in platycalycal type and Glossograptidae with two stipes side to side (monopleural) developed in pericalycal type. Morphologically, *Oncograptus* is an intermediate form between *Isograptus* and *Cardigraptus*. Two distinct series in *Oncograptus* and *Cardigraptus* are recognized, namely, the *upsilon-morsus* series with wedge-shaped rhabdosome and the *magnus-amplus* series with parallel-sided rhabdosome. The former occurs in North America, Australasia and the NW Region of China, whereas the latter is only known in the S. China Region. The relation between *Oncograptus magnus* and *Cardigraptus amplus* are possibly transitional, whereas the relation between *Oncograptus upsilon* and *Cardigraptus morsus* is uncertain, because an additional theca is present in *Oncograptus upsilon* (Bulman, 1936) and a distinct virgella is present in an allied species *Oncograptus zhongguensis* Xu & Huang (1979). Therefore, *Oncograptus*

graptus is here placed in Isograptidae of Axonolipa. Skevington (1968) considered *Oncograptus upsilon* and *Cardigraptus morsus* to be conspecific derived from "*Isograptus*" *manubriatus* based on the so-called "curved" proximal thecae. Cooper (1979) demonstrated the proximal thecae in *Oncograptus* and *Cardigraptus* which grow downwards without curvature.

The characteristic feature of Glossograptidae is the overlapping of the two scandent stipes (monopleural). In *Apiograptus* Cooper & McLaurin (1974) the two stipes begin to be overlapped laterally (Cooper, 1979). It was considered to be a primitive member of Glossograptidae (= Cryptograptidae) by Mu et al., 1979. In the advanced form, *Cryptograptus tricornis*, the two stipes are entirely overlapped. For the sake of clarity, the writer used formerly Hadding's Cryptograptidae instead of Lapworth's Glossograptidae, because Lapworth's Glossograptidae is fused with his Lasiograptidae. For the same reason, the writer proposed Hallograptidae instead of Lapworth's Lasiograptidae (Mu, 1950; Urbanek, 1959). It is unnecessary to divide the family Glossograptidae (= Cryptograptidae) into two families (Strachan, 1985).

Anagenetic Grades and Cladogenetic Divergences

As stated above, Axonocrypta links with Axonolipa and Axonophora. Dendrograptidae and Anisograptidae of Dendroidea, Tetragraptidae-Didymograptidae and Isograptidae of Axonolipa, Cardigraptidae of Axonocrypta and Diplograptidae and Monograptidae of Axonophora represent anagenetic grades. Some important evolutionary trends such as simplification and complication of rhabdosome, overlapping of stipes, elaboration of thecae and reduction of periderm, took place once again in the graptolite history representing cladogenetic divergences. The following taxa are all offshoots of various grades.

Simplification of rhabdosome: Azygograptidae and Corynoididae in Axonolipa, Peiragraptidae and Dimorphograptidae in Axonophora.

Complication of rhabdosome: Pterograptinae, Nemagraptinae (= Pleurograptinae) and Tangyagraptinae in Axonolipa, Diversograptinae,

Cyrtograptinae and Linograptinae in Axonophora.

Overlapping of stipes: Kalpinograptidae in Axonolipa, Glossograptidae (= Cryptograptidae) in Axonocrypta.

Elaboration of thecae: Muenzhingraptidae-Psiograptidae in Dendroidea, Sinograptidae, Kinnegraptidae, Atopograptidae and Dicranograptidae in Axonolipa, various lineages in diplograptids and monograptids in Axonophora.

Reduction of periderm: Abrograptidae in Axonolipa, Reteograptidae and Archiretiolitidae-Retiolitidae-Plectograptidae in Axonophora.

Studies on internal structure of graptolites afford important features of taxonomic significance:

1) The initial bud growing from the porus on the prosicula or on the proximal part of metasicula changes downwards to grow from the distal part of metasicula and the porus changes to a sinus.

2) The formation of a virgella beginning at the distal part of metasicula changes upwards to be at the proximal part of the metasicula.

3) The growth direction of the proximal thecae especially the first two pairs changes from downwards to upwards.

4) The formation of the median septum in scandent rhabdosome changes from a compound septum (median septa in Axonocrypta) to a simple septum (in Axonophora).

Ultrastructural studies of graptolites have provided some important information for graptolite taxonomy as reviewed by Rickards, Crowther & Chapman (1982). Further studies on the microstructure and ultrastructure of the etched materials will settle well the problem of the graptolite taxonomy.

Dansk sammendrag

Graptolithina er sammensat af seks ordener. Blandt dem er Graptoloidea og en del af Dendroidea kendt for deres planktoniske levevis. Graptoloidea er sammensat af tre underordener nemlig Axonolipa, Axonocrypta og Axonophora. Familierne Dendrograptidae-Anisograptidae-Tetragraptidae og Didymograptidae-Isograptidae-Cardigraptidae-Diplograptidae-Monograptidae repræsenterer anagenetiske grader. Nogle vigtige evolutionære tendenser viste sig igen i form af cladogenetiske afspaltninger. Alle andre familier eller subfamilier er sideskud af forskellig grad.

Underordenen Axonocrypta bliver diskuteret i nogen detalje i det nærværende arbejde.

References

- Bjerreskov, M., 1975: Llandoveryan and Wenlockian graptolites from Bornholm, *Fossils & Strata*, 8., 94 pp.
- Bulman, O. M. B., 1936: The structure of *Oncograptus* T. S. Hall, *Geol. Mag.*, 73, 864, 271–278.
- Bulman, O. M. B., 1963: The evolution and classification of the graptoloidea, *Quart. Jl. geol. Soc. London*, 119, 401–418.
- Bulman, O. M. B., 1970: Graptolithina with section on Enteropneusta and Pterobranchia, *Treatise on Invertebrate Paleontology*, part V, Geol. Soc. Amer. and Univ. Kansas, 163 pp.
- Chen Xu, 1982, Early Ordovician *Exigraptus* and *Glyptograptus* from Xingan, N. Guangxi and the origin of Biseriis Axonophorous graptoloids, *Acta Pal. Sinica*, 21, 5, 513–535.
- Cooper, R. A., 1979: Ordovician geology and graptolite faunas of the Aorangi Mine area, Northwest Nelson, New Zealand, *Pal. Bull.*, 47, New Zealand Geological Survey, 1–127.
- Cooper, R. A. & Fortey, R. A., 1982: The Ordovician Graptolites of Spitsbergen, *Bull. Br. Mus. Nat. Hist. (Geol.)* 36, 3, 157–302.
- Cooper, R. A. & Fortey, R. A., 1983: Development of the graptoloid rhabdosome, *Alcheringa*, 7, 201–221.
- Cooper, R. A. and Lindholm, K., 1985: The phylogenetic relationships of the graptolites *Tetragraptus phyllograptoides* and *Pseudophyllograptus cor.* *Geol. Foren. Stockholm Forh.* 106, 3, 279–291.
- Cooper, R. A. and McLaurin, A. N., 1974: *Apiograptus* gen. nov. and the origin of the biserial graptoloid rhabdosome, *Special papers in Palaeontology*, 13, 75–85.
- Finney, S. C., 1978: The affinities of *Isograptus*, *Glossograptus*, *Cryptograptus*, *Corynoides*, and allied graptolites. *Acta Palaeontologica Polonica*, 23, 4, 481–495.
- Finney, S. C. 1980: Thamnograptid, Dichograptid and Abrograptid graptolites from the Middle Ordovician Athens Shale of Alabama, *Jour. Palaeont.*, 54, 6, 1184–1209.
- Fortey, R. A., 1971: *Tristichograptus*, A Triserial Graptolite from the Lower Ordovician of Spitsbergen, *Palaeontology*, 14, 1, 188–199.
- Frech, F., 1897: *Lethaea geognostica* I, *Lethaea palaeozoica*, 1, Leipzig, 690 pp.
- Hsu, S. C., 1934: Graptolites of the Lower Yangtze Valley, *Monograph Nat. Res. Inst. Geol.*, A 4, 106 pp.
- Hsu, S. C. & Chen, P. L., 1964: On the genus *Trigonograptus*, *Scientia Sinica*, 13, 4, 639–653.
- Jaanusson, V., 1960: Graptoloids from the Ontikan and Viruan (Ordov.) Limestones of Estonia and Sweden, *Bull. geol. Inst. Univ. Uppsala*, 38, 289–366.
- Jenkins, C., 1980: *Maeandrograptus schmalenseeii* and its bearing on the origin of the diplograptids. *Lethaia*, 13, 289–302.
- Kozłowski, R., 1949: Les Graptolithes et Quelques Nouveaux Groupes d'Animaux du Tremadoc de la Pologne. *Paleontologia Polonica*, iii, 1948, 235 pp.
- Kozłowski, R., 1954: Sur la structure de certains Dichograptids. *Acta Geol. Polonica*, 4, 423–444.
- Kozłowski, R., 1966: On the structure and relationships of graptolites, *Jour. Palaeont.* 40, 3, 489–501.
- Lapworth, C., 1980: On the geological distribution of the Rhabdophora. *Ann. Mag. Nat. Hist.* V, 5, 45–62, 273–285, 359–369; vol. 6, pp. 16–29, 185–207.
- Lin Yao-kun, 1981: New Materials of Graptodendroids with special reference to the Graptodendroidea. *Bull. Nanjing Inst. Geol. Paleont. Acad. Sinica*, 3, 241–262.
- Mu, A. T., 1950: On the evolution and classification of Graptoloids. *Ti-chih-lun-ping*, 15, 4–6, 171–183. (in Chinese).
- Mu, A. T., 1974: Evolution, Classification and Distribution of Graptoloidea and Graptodendroids, *Scientia Sinica*, 17, 2, 227–238.
- Mu, A. T. & Lee, C. K., 1958: Scandent graptolites from the Nankuo Shale of Kiangshan-Changshan area, western Chekiang, *Acta Palaeont. Sin.*, 6, 4, 391–428.
- Mu, A. T., Lee, C. K. & Geh, M. Y., 1960: Ordovician graptolites from Xinjiang, *Acta Palaeont. Sinica*, 8, 1, 27–39.
- Mu, A. T. & Zhan, S. G., 1966: On the probable development and systematic position of *Glossograptus*. *Scientia Sinica*, 15, 92–98.
- Mu En-zhi, Ge Mei-yu, Chen Xu, Ni Yu-nan & Lin Yao-kun, 1979, Lower Ordovician graptolites of wouthwest China. *Palaeont. Sinica*, 156B (13), 1–1921.
- Obut, A. M., 1957: Classification and ordering of the genera of graptolites. *Vsesoyuz. Paleont. Obshch. Ezhegodnik* 16, 11–47. (in Russian).
- Rickards, R. B., 1973: The Arenig graptolite genus *Pseudotrigonograptus* Mu et Lee, 1958, *Acta geol. Polonica*, 23, 4, 597–604.
- Rickards, R. B., 1974: A new monograptid genus and the origins of the main monograptid genera. *Special Paper in Palaeontology* 13, 141–147.
- Rickards, R. B., Crowther, P. R. and Chapman, A. J., 1982, Ultrastructural studies of graptolites – a review, *Geol. Mag.*, 119, 4, 355–370.
- Ruedemann, R., 1904: Graptolites of New York, Part I: Graptolites of the lower beds. *N. Y. State Mus., Mem.*, 7, 457–803.
- Ruedemann, R., 1908: Graptolites of New York, part II: Graptolites of higher beds. *N. Y. State Mus., Mem.* 11, 1–583.
- Skevington, D., 1968: The affinities of *Oncograptus*, *Cardiograptus* and allied graptolites from the Lower Ordovician, *Lethaia*, 1, 4, 311–324.
- Strachan, I., 1985: The significance of the proximal end of *Cryptograptus tricornis* (Carruthers) (Graptolithina), *Geol. Mag.* 122, 2, 151–155.
- Urbanek, A., 1959: Studies on graptolites II, on the development and structure of graptolite genus *Gymnograptus* Bulman, *Acta Palaeont. Polonica*, IV, 3, 279–338.
- Xu Jie & Huang Zhi-gao, 1979: Lower Ordovician graptolite faunas of Guozigou Area, Hocheng, Xinjiang. *Acta geol. Sinica*, 1979, 1, 1–19.
- Yu Jian-hua & Fang Yi-ting, 1979: On the classification of graptoloids. *Acta Palaeont. Sinica*, 18, 5, 435–443.
- Zhao Xiang-lin & Shang Zhun-xin, 1985: Reclined graptolites of the Xinchangian. *Journal of Changchun College of Geology*, 1985 (2), 13–26.